
## U.S. study suggests COVID is less severe with Omicron than Delta



The Omicron variant appears to result in less severe COVID-19 than seen during previous periods of high coronavirus transmission including the Delta wave, with shorter hospital stays, less need for intensive care and fewer deaths, according to a new U.S. study.

Atlanta, January 27 (RHC)-- The Omicron variant appears to result in less severe COVID-19 than seen during previous periods of high coronavirus transmission including the Delta wave, with shorter hospital stays, less need for intensive care and fewer deaths, according to a new U.S. study.

However, the fast-spreading Omicron variant has led to record numbers of infections and hospitalizations, straining the U.S. healthcare system.

Despite the steep spike in COVID cases, the percentage of hospitalized patients admitted to intensive care units (ICU) during the current Omicron wave was about 29% lower than during last winter's surge and some 26% lower than during the Delta wave, the study published in the U.S. Centers for Disease Control and Prevention's (CDC) Morbidity and Mortality Weekly Report found.

The lower COVID-19 disease severity during the Omicron period is likely related to higher vaccination coverage, booster use among those eligible for the extra shots, as well as prior infections providing some immune protection, the study said.

Deaths in the period from December 19 to January 15, when Omicron infections were at a peak, averaged 9 per 1,000 COVID cases, compared to 16 per 1,000 in the previous winter peak and 13 during the Delta wave, the study showed.

The findings were consistent with previous data analyses from South Africa, England and Scotland, where infections from Omicron peaked earlier than in the United States, the CDC said. Relatively high hospitalizations among children during the Omicron period may be related to lower vaccination rates compared with adults, the agency said. Children under age 5 are not yet eligible for vaccines in the United States and the rate of vaccination among older children lags that of adults.

The study involved analysis of data from a large healthcare database and three surveillance systems to assess U.S. COVID-19 characteristics from December 1, 2020 to January 15, 2022.

The authors said one limitation of the study was that it was unable to exclude incidental infections in which patients admitted for other reasons test positive for COVID while in the hospital. That may inflate hospitalization-to-case ratios and affect severity indicators.

 $\frac{https://www.radiohc.cu/en/noticias/internacionales/284273-us-study-suggests-covid-is-less-severe-with-omicron-than-delta}{omicron-than-delta}$ 



Radio Habana Cuba